arXiv:1105.5530v1 [math-ph] 27 May 2011

A REMARK ON EXACT FORMULAS FOR THE RIESZ ENERGY OF THE NTH ROOTS OF UNITY

J. S. BRAUCHART

Abstract. In the paper the author, Hardin and Saff [Bull. Lond. Math. Soc. 41 (2009), no. 4] we provided the complete asymptotic expansions of the Riesz s-energy of the Nth roots of unity. In this note the exact formulas (valid for all $N \geq 2$) are obtained for s is an even integer. Explicit representations of the coefficients of the powers of N are derived by direct computations. In the case of positive s a continuous modified energy approximation of the Riesz energy is used. Stirling numbers of the first kind, Eulerian numbers and special values of partial Bell polynomials play a central role. A number of identities between these quantities are shown.

The Riesz s-energy (real $s \neq 0$) of an N-point configuration $X_N = \{z_1, \dots, z_N\}$ in the complex plane (identified with the Euclidean space $\mathbb{R}^2$) is given by

$$\mathrm{E}_s(X_N) = \sum_{\substack{j=1 \\ j \neq k}}^{N} \sum_{k=1}^{N} \frac{1}{\left| z_j - z_k \right|^s}.$$

It can be shown (see [4]) that the Nth roots of unity (or any rotation thereof) maximize (if $-2 < s < 0$) and minimize (if $s > 0$) the above energy functional over all N-point configuration on the unit circle. We remark that for $s < -2$ an optimal configuration on the unit circle consists of half of the points at one of two opposite locations. (This follows from results of G. Björck [3].)

The Riesz s-energy (if $s \neq 0$ is real) of the N-th roots of unity can be evaluated by means of

$$\mathcal{L}_s(N) = 2^{-s} N \sum_{k=1}^{N-1} \left(\sin \frac{\pi k}{N} \right)^{-s}, \qquad N \geq 2, \, s \in \mathbb{C}, \, s \neq 0. \tag{1}$$

It has the following general complete asymptotic expansion of $\mathcal{L}_s(N)$ ([4, Thm. 1.1])

$$\mathcal{L}_s(N) = V_s \, N^2 + \frac{2}{(2\pi)^s} \sum_{n=0}^{p} \alpha_n(s) \, \zeta(s-2n) N^{1+s-2n} + \mathcal{O}_{s,p}(N^{-1+\mathrm{Re}\, s - 2p}), \qquad \text{as } N \to \infty, \tag{2}$$

provided s is not a positive odd integer. (These exceptional cases are considered in [4, Thm. 1.2].)

The Riemann zeta function has its trivial zeros at negative even integers. Thus, the sum in (2) vanishes for all $N \geq 2$ if s is a negative even integer or can have at most m terms if $s = 2m$ is a positive even integer.

The case $s = -2m$ and m a positive integer. In [4, Remark 1, Eq. (1.19)] we observed that

$$\mathcal{L}_s(N) = V_s N^2, \qquad s = -2, -4, -6, \dots,$$

V_s given in (4), which is **correct up to a remainder term of order $\mathcal{O}_{s,p}(N^{-1-s-2p})$, where p can be any positive integer**. The precise result is as follows.

Proposition 1. *Let m be a positive integer and set $s = -2m$. Then*

$$\mathcal{L}_s(N) = V_s \, N^2 + 2N^2 \sum_{\substack{k=1 \\ N \mid k}}^{m} (-1)^k \binom{2m}{m-k}, \qquad N = 2, 3, 4, \dots,$$

where $N|k$ denotes that the integer N divides the integer k. The sum above vanishes if $N > m$.

Date: January 26, 2013.

Key words and phrases. Bell polynomials, Eulerian numbers, Riesz energy, Stirling numbers.

For the sake of completeness we include here the proof.

Proof. Using the identity ([8, Eq. I.1.9])

$$(\sin\phi)^{2m} = 2^{1-2m}\sum_{k=0}^{m-1}(-1)^{m-k}\binom{2m}{k}\cos[2(m-k)\phi] + 2^{-2m}\binom{2m}{m},$$

in (1) we get

$$\mathcal{L}_{-2m}(N) = 2^{2m}N\sum_{k=0}^{N-1}\left(\sin\frac{\pi k}{N}\right)^{2m} = 2N\sum_{\ell=1}^{m}(-1)^{\ell}\binom{2m}{m-\ell}\left\{\sum_{k=0}^{N-1}\cos\frac{2\pi k\ell}{N}\right\} + \binom{2m}{m}N^2$$

a representation in terms of trigonometric sums satisfying the identity ([8, Eq.s 4.4.4.6–8])

$$\sum_{k=0}^{N-1}\cos(2\pi k\ell/N) = \begin{cases} N & \text{if } N \text{ divides } \ell \text{ (that is } N|\ell\text{)},\\ 0 & \text{otherwise.}\end{cases} \tag{3}$$

Also observe that [4, Eq. (1.8)] can be written as

$$\begin{aligned} V_s &= 2^{-s}\frac{\Gamma((1-s)/2)}{\sqrt{\pi}\,\Gamma(1-s/2)} = 2^{2m}\frac{\Gamma((1+2m)/2)}{\sqrt{\pi}\,\Gamma(1+m)} = 2^{2m}\frac{\Gamma(m+1/2)\,\Gamma(1+m)}{\sqrt{\pi}\,\Gamma(1+m)\,\Gamma(1+m)} \\ &= \frac{\Gamma(2m+1)}{\Gamma(m+1)\,\Gamma(m+1)} = \binom{2m}{m} \end{aligned} \tag{4}$$

by the duplication formula for the gamma function. The result follows. □

The case $s = 2m$ and m a positive integer. In [4, Remark 1, Eq. (1.20)] it was observed that

$$\mathcal{L}_s(N) = \frac{2}{(2\pi)^s}\sum_{n=0}^{m}\alpha_n(s)\,\zeta(s-2n)N^{1+s-2n}, \qquad s = 2m, m = 1, 2, 3, \dots, \tag{5}$$

by arguing that in the general asymptotic expansion (2) the constant V_s given in (4) vanishes if s is a positive even integer and the Riemann zeta function part $\zeta(s-2n)$ assumes the value 0 for $s-2n$ is an even negative integer. One should add that (5) is true **up to a remainder term of order $\mathcal{O}_{s,p}(N^{-1-s-2p})$, where p can be any positive integer**. In fact, direct computations suggest that relation (5) holds for all $N \geq 2$. The coefficients $\alpha_n(s)$ are defined by the following generating function relation and can be expressed explicitly in terms of *generalized Bernoulli polynomials* $B_k^{(\sigma)}(x)$; that is

$$\left(\frac{\sin\pi z}{\pi z}\right)^{-s} = \sum_{n=0}^{\infty}\alpha_n(s)z^{2n}, \quad |z|<1,\ s\in\mathbb{C}, \qquad \alpha_n(s) = \frac{(-1)^n B_{2n}^{(s)}(s/2)}{(2n)!}(2\pi)^{2n}, \quad n \geq 0. \tag{6}$$

For non negative even integers the Riemann zeta function can be expressed in terms of Bernoulli numbers by means of $\zeta(2m) = 2^{2m-1}(-1)^{m-1}B_{2m}\pi^{2m}/(2m)!$, $m = 0, 1, 2, \dots$.

The validity of relation (5) for all $N \geq 2$ follows by means of a limit process from the corresponding result for a modified (and thus non-singular) form of the Riesz energy of the Nth roots of unity.

The distance between a point $\mathbf{x}$ on the unit circle ($|\mathbf{x}| = 1$) and a point $\mathbf{y}$ inside the unit circle ($r = |\mathbf{y}| < 1$) is given by

$$|\mathbf{x}-\mathbf{y}|^2 = 1 - 2r\langle\mathbf{x},\mathbf{y}\rangle + r^2 = (1-r)^2 + 4r\left[\sin(\phi/2)\right]^2, \qquad \cos\phi = \langle\mathbf{x},\mathbf{y}\rangle.$$

This motivates the following modification of the the s-energy of the N-th roots of unity:

$$\mathcal{M}_s(N;r) := N\sum_{k=1}^{N-1}\left(1 - 2r\cos\frac{2\pi k}{N} + r^2\right)^{-s/2}, \qquad 0 \leq r < 1. \tag{7}$$

Observe that $\mathcal{L}_s(N) = \lim_{r\to 1^-} \mathcal{M}_s(N;r)$. It should be mentioned that Chu [5] considered sums of the form

$$f_m(z) = \sum_{k=0}^{n-1} \frac{1-z^2}{\left[1-2z\cos(2\pi k/n)+z^2\right]^m},$$

where m and n are positive integers and z is complex. The following recurrence relation is given:

$$f_{m+1}(z) = \frac{f_m(z)}{1-z^2} + \frac{z}{m}\frac{\mathrm{d}}{\mathrm{d}\,z}\left\{\frac{f_m(z)}{1-z^2}\right\}.$$

Chu's main interest was in the case when $z=\sqrt{-1}$.

We make use of the *Gauss hypergeometric function*

$$ {}_2\mathrm{F}_1\left({a,b \atop c};z\right) = \sum_{n=0}^{\infty} \frac{(a)_n(b)_n}{(c)_n}\frac{z^n}{n!}, \qquad |z|<1, \tag{8}$$

where $(a)_n = \Gamma(n+a)/\Gamma(a)$ denotes the Pochhammer symbol.

Proposition 2. *Let $s>0$. Then for integers $N \geq 2$*

$$\mathcal{M}_s(N;r) = N^2\, G_0(s;r) - N\,(1-r)^{-s} + 2N^2 \sum_{\nu=1}^{\infty} G_{\nu N}(s;r), \qquad 0\leq r<1,$$

where the series is uniformly convergent with respect to r on compact subsets of $[0,1)$ and

$$G_n(s;r) = \left(1-r^2\right)^{1-s}\frac{(s/2)_n}{n!} r^n\, {}_2\mathrm{F}_1\left({1-s/2, n+1-s/2 \atop n+1};r^2\right).$$

For s a positive even integer, that is $s=2m$, we can write (cf. [1, Eq. 15.8.7])

$$G_n(2m;r) = \left(1-r^2\right)^{1-2m}\frac{2^{2m-2}\,\Gamma(m-1/2)}{\sqrt{\pi}\,\Gamma(m)} r^n\, {}_2\mathrm{F}_1\left({1-m, n+1-m \atop 2-2m};1-r^2\right). \tag{9}$$

The hypergeometric function above reduces to a polynomial of degree $m-1$.

For $s=2$, that is $m=1$, one simply has $G_n(2;r) = r^n/(1-r^2)$ which leads to

$$\begin{aligned}\mathcal{M}_2(N;r) &= \frac{N^2}{1-r^2} - \frac{N}{(1-r)^2} + \frac{2N^2}{1-r^2}\sum_{\nu=1}^{\infty} r^{\nu N} = \frac{N^2}{1-r^2} - \frac{N}{(1-r)^2} + \frac{2N^2 r^N}{(1-r^2)\,(1-r^N)} \\ &= \frac{1+r^N}{(1-r^2)\,(1-r^N)}\,N^2 - \frac{N}{(1-r)^2}.\end{aligned}$$

Using Mathematica, a limit process $r\to 1^-$ yields that

$$\mathcal{L}_2(N) = \lim_{r\to 1^-}\mathcal{M}_2(N;r) = \frac{N^3}{12} - \frac{N}{12} \qquad \text{and by (5):} \qquad \mathcal{L}_2(N) = \frac{2}{4\pi^2}\left(\frac{\pi^2}{6}N^3 - \frac{\pi^2}{6}N\right).$$

It follows that (5) is correct for $s=2$ and all $N\geq 1$.

In the general result we make use of the *Stirling numbers of the first kind* $\mathrm{s}(n,k)$ defined by the expansion ([1, Eq. 26.8.7])

$$\sum_{k=0}^{n} \mathrm{s}(n,k)x^k = x\,(x-1)\cdots(x-n+1) = (x+1-n)_n, \tag{10}$$

where $(a)_n$ is the Pochhammer symbol, and the *Eulerian numbers* $\left\langle {n \atop k} \right\rangle$ given by (cf. [1, Eq.s 26.14.5 and 26.14.6])

$$\sum_{k=0}^{n-1}\left\langle {n \atop k}\right\rangle \binom{x+k}{n} = x^n, \qquad \left\langle {n \atop k}\right\rangle = \sum_{j=0}^{k}(-1)^j\binom{n+1}{j}(k+1-j)^n \qquad n\geq 1. \tag{11}$$

There holds $\left\langle {0 \atop 0}\right\rangle = \left\langle {n \atop 0}\right\rangle = 1$ and $\left\langle {n \atop n-1}\right\rangle = 1$. By convention we set $\left\langle {n \atop k}\right\rangle = 0$ for $k\geq n$.

Proposition 3. *Let* m *and* N *be positive integers. Then*

$$\mathcal{M}_{2m}(N;r) = \frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} \left[-(1-m)_k + 2 \sum_{q=0}^{k} g(k,q;N,1-m) \left(1-r^N\right)^{-q-1} \right] \left(1-r^2\right)^{k+1-2m} - N\left(1-r\right)^{-2m}, \qquad 0 \leq r < 1,$$

where

$$g(k,q;a,b) = \sum_{p=0}^{k-q} (-1)^p s(k,p+q;b) b(p+q,p) a^{p+q}, \qquad b(p+q,p) = \sum_{j=0}^{q} \left\langle {p+q \atop j} \right\rangle \binom{p+q-j}{p},$$

$$\mathrm{s}(n,\ell;y) = \sum_{k=\ell}^{n} \binom{k}{\ell} \mathrm{s}(n,k) \left(y+n-1\right)^{k-\ell}.$$

We are interested in the limit as $r \to 1^-$. Clearly, $\mathcal{M}_{2m}(N) := \lim_{r\to 1^-} \mathcal{M}_{2m}(N;r) = \mathcal{L}_{2m}(N)$.

Proposition 4. *Let* m *and* N *be positive integers. Then*

$$\mathcal{M}_2(N) = \frac{N^3}{12} - \frac{N}{12}, \qquad \mathcal{M}_{2m}(N) = \sum_{\nu=0}^{2m} \beta_{2m-\nu}(m)\, N^{1+2m-\nu}, \quad m \geq 2, \tag{12}$$

where the coefficients $\beta_\nu(m)$ $(0 \leq \nu \leq 2m)$ *do not depend on* N.

Explicit expressions for the coefficients $\beta_\nu(m)$ are given in the Proof of Prop. 4 (see Eq.s (19)). The expansion (12) holds for every integer $N \geq 2$, whereas the expansion (5) (derived from an asymptotic result as $N \to \infty$) representing the same quantity holds for sufficiently large N. Since the coefficients of the powers of N in either expansion do not depend on N, by comparing (19) and (5) we obtain for $s = 2m$ and m a positive integer the connection formulas

$$\beta_{2n+1}(m) = 0 \quad (0 \leq n < m), \qquad \beta_{2m-2n}(m) = \frac{2\,\zeta(s-2n)}{(2\pi)^s} \alpha_n(s) \quad (0 \leq n \leq m).$$

The coefficients $\alpha_n(2m)$ can be expressed in terms of generalized Bernoulli polynomials and the Riemann zeta function for nonnegative even integers can be expressed in terms of Bernoulli numbers (cf. (6) and text there). This leads to

$$\beta_{2m-2n}(m) = \frac{(-1)^{m-n-1} B_{2m-2n}}{(2m-2n)!} \frac{(-1)^n B_{2n}^{(2m)}(m)}{(2n)!}, \qquad 0 \leq n \leq m.$$

For $n = 0$ one gets the following identity.

Proposition 5. *For* $m = 1, 2, \dots$ *on has*

$$\frac{(-1)^{m-1} B_{2m}}{(2m)!} = \sum_{\nu=0}^{m-1} \sum_{p=0}^{\nu} (-1)^p \frac{2^{2+\nu-2m}\, \Gamma(2m-1-\nu)}{\Gamma(m)\nu!\, \Gamma(m-\nu)} b(\nu,p) G(2m-p, 2m-p, \nu-p+1),$$

where the numbers $b(\nu,p)$ *and* $G(n,n,q)$ *are given by*

$$b(k,p) = \sum_{j=0}^{k-p} \left\langle {k \atop j} \right\rangle \binom{k-j}{p}, \qquad \sum_{n=1}^{\infty} (-1)^n G(n,n,q)\, x^n = \left(\frac{e^x - 1}{x} \right)^{-q} - 1.$$

1. Proofs

The Gegenbauer polynomials $C_n^{(\lambda)}$ of degree n with index $\alpha > 0$ may be defined by means of the generating function relation

$$\sum_{n=0}^{\infty} z^n\, C_n^{(\lambda)}(\cos\phi) = \frac{1}{\left(1 - 2z\cos\phi + z^2\right)^{\lambda}}, \qquad |z| < 1, \lambda > 0. \tag{13}$$

They admit a representation through trigonometric functions (cf. [2, Section 22]) as follows

$$C_n^{(\lambda)}(\cos\phi) = \sum_{\ell=0}^{n} \frac{(\lambda)_\ell (\lambda)_{n-\ell}}{\ell!(n-\ell)!} \cos[(n-2\ell)\phi], \qquad \lambda \neq 0. \tag{14}$$

We need the following auxiliary result.

Lemma 6. *Let $s > 0$. Then*

$$\mathcal{A}_n(s;N) := N \sum_{k=1}^{N-1} C_n^{(s/2)}(\cos\frac{2\pi k}{N}) = N^2 \sum_{\substack{\ell=0\\ N|(n-2\ell)}}^{n} \frac{(s/2)_\ell (s/2)_{n-\ell}}{\ell!(n-\ell)!} - N\, C_n^{(s/2)}(1).$$

For even and odd n one has

$$\mathcal{A}_{2\nu}(s;N) = \frac{(s/2)_\nu (s/2)_\nu}{\nu!\nu!} N^2 - C_{2\nu}^{(s/2)}(1)\, N + 2N^2 \sum_{\substack{\ell=1\\ N|(2\ell)}}^{\nu} \frac{(s/2)_{\nu-\ell}(s/2)_{\nu+\ell}}{(\nu-\ell)!(\nu+\ell)!},$$

$$\mathcal{A}_{2\nu+1}(s;N) = 2N^2 \sum_{\substack{\ell=0\\ N|(2\ell+1)}}^{\nu} \frac{(s/2)_{\nu-\ell}(s/2)_{\nu+1+\ell}}{(\nu-\ell)!(\nu+1+\ell)!} - C_{2\nu+1}^{(s/2)}(1)\, N.$$

Proof. By (14) and (3) we have

$$\mathcal{A}_n(s;N) = \sum_{\ell=0}^{n} \frac{(s/2)_\ell (s/2)_{n-\ell}}{\ell!(n-\ell)!} N \sum_{k=1}^{N-1} \cos[(n-2\ell)\frac{2\pi k}{N}] = N^2 \sum_{\substack{\ell=0\\ N|(n-2\ell)}}^{n} \frac{(s/2)_\ell (s/2)_{n-\ell}}{\ell!(n-\ell)!} - N\, C_n^{(s/2)}(1).$$

Since $N|(n-2\ell)$ if and only if $N|(2\ell-n)$, where $2\ell-n=-[n-2(n-\ell)]$, we can use symmetry to write

$$\sum_{\substack{\ell=0\\ N|(2\nu-2\ell)}}^{2\nu} \frac{(s/2)_\ell (s/2)_{2\nu-\ell}}{\ell!(2\nu-\ell)!} = 2 \sum_{\substack{\ell=0\\ N|2(\nu-\ell)}}^{\nu-1} \frac{(s/2)_\ell (s/2)_{2\nu-\ell}}{\ell!(2\nu-\ell)!} + \frac{(s/2)_\nu (s/2)_\nu}{\nu!\nu!},$$

$$\sum_{\substack{\ell=0\\ N|(2\nu+1-2\ell)}}^{2\nu+1} \frac{(s/2)_\ell (s/2)_{2\nu+1-\ell}}{\ell!(2\nu+1-\ell)!} = 2 \sum_{\substack{\ell=0\\ N|(2\nu+1-2\ell)}}^{\nu} \frac{(s/2)_\ell (s/2)_{2\nu+1-\ell}}{\ell!(2\nu+1-\ell)!}.$$

The results for even and odd n follow by reversing the order of terms on the right-hand sides above and substituting the expressions into the result for general n. □

Proof of Proposition 2. Using (13) in (7) and Lemma 6, we obtain

$$\mathcal{M}_s(N;r) = \sum_{n=0}^{\infty} r^n\, N \sum_{k=1}^{N-1} C_n^{(s/2)}(\cos\frac{2\pi k}{N}) = N^2 \sum_{n=0}^{\infty} r^n \sum_{\substack{\ell=0\\ N|(n-2\ell)}}^{n} \frac{(s/2)_\ell (s/2)_{n-\ell}}{\ell!(n-\ell)!} - N \sum_{n=0}^{\infty} r^n\, C_n^{(s/2)}(1).$$

The last series has the closed form $(1-r)^{-s}$ (cf. (13)). One has

$$\begin{aligned}
\sum_{\nu=0}^{\infty} r^{2\nu} \sum_{\substack{\ell=0\\ N|(2\nu-2\ell)}}^{2\nu} \frac{(s/2)_\ell (s/2)_{2\nu-\ell}}{\ell!(2\nu-\ell)!} &= \sum_{\nu=0}^{\infty} \frac{(s/2)_\nu (s/2)_\nu}{\nu!\nu!} r^{2\nu} + 2\sum_{\nu=0}^{\infty} r^{2\nu} \sum_{\substack{\ell=1\\ N|(2\ell)}}^{\nu} \frac{(s/2)_{\nu-\ell}(s/2)_{\nu+\ell}}{(\nu-\ell)!(\nu+\ell)!} \\
&= \sum_{\nu=0}^{\infty} \frac{(s/2)_\nu (s/2)_\nu}{\nu!\nu!} r^{2\nu} + 2 \sum_{\substack{\ell=1\\ N|(2\ell)}}^{\infty} \sum_{\nu=\ell}^{\infty} \frac{(s/2)_{\nu-\ell}(s/2)_{\nu+\ell}}{(\nu-\ell)!(\nu+\ell)!} r^{2\nu} \\
&= {}_2\mathrm{F}_1\Big(\begin{matrix} s/2, s/2 \\ 1 \end{matrix}; r^2\Big) + 2 \sum_{\substack{\ell=1\\ N|(2\ell)}}^{\infty} \frac{(s/2)_{2\ell}}{(2\ell)!} r^{2\ell}\, {}_2\mathrm{F}_1\Big(\begin{matrix} s/2, 2\ell+s/2 \\ 2\ell+1 \end{matrix}; r^2\Big).
\end{aligned}$$

In the last step (8) was used. By the same token

$$\sum_{\nu=0}^{\infty} r^{2\nu+1} \sum_{\substack{\ell=0\\ N\mid(2\nu+1-2\ell)}}^{2\nu} \frac{(s/2)_\ell (s/2)_{2\nu+1-\ell}}{\ell!(2\nu+1-\ell)!} = 2 \sum_{\substack{\ell=1\\ N\mid(2\ell+1)}}^{\infty} \frac{(s/2)_{2\ell+1}}{(2\ell+1)!} r^{2\ell+1}\, {}_2\mathrm{F}_1\left(\begin{matrix} s/2, 2\ell+1+s/2\\ 2\ell+2\end{matrix}; r^2\right).$$

Putting everything together we obtain

$$\mathcal{M}_s(N;r) = N^2\, G_0(s;r) - N\,(1-r)^{-s} + 2N^2 \sum_{\substack{n=1\\ N\mid n}}^{\infty} G_n(s;r), \qquad G_n(s;r) := \frac{(s/2)_n}{n!} r^n\, {}_2\mathrm{F}_1\left(\begin{matrix} s/2, n+s/2\\ n+1\end{matrix}; r^2\right).$$

The series $\sum_{\nu=1}^{\infty} G_{\nu N}(s;r)$ converges uniformly with respect to r on compact subsets of $[0,1)$, which can be seen from the integral representation (cf. [1, Eq. 15.6.1]) and estimate

$$\begin{aligned} G_n(s;r) &= r^n \frac{1}{[\Gamma(s/2)]^2} \frac{\Gamma(n+s/2)}{\Gamma(n+1-s/2)} \int_0^1 \frac{t^{s/2-1}\,(1-t)^{n-s/2}}{(1-r^2t)^{n+s/2}}\, \mathrm{d}\, t \\ &\leq c_s(r) \frac{\Gamma(n+s/2)}{\Gamma(n+1-s/2)}\, r^n, \qquad c_s(r) = \frac{1}{[\Gamma(s/2)]^2} \int_0^1 \frac{t^{s/2-1}}{(1-r^2t)^s}\, \mathrm{d}\, t, \end{aligned}$$

valid for $n \geq n_0 > s/2$, where $c_s(r)$ can be uniformly bounded on compact sets in $[0,1)$. Hence

$$0 < \sum_{\nu=1}^{\infty} G_{\nu N}(s;r) \leq C + \sum_{n=n_0}^{\infty} G_n(s;r) \leq C + c_s(r) \sum_{n=n_0}^{\infty} \frac{\Gamma(n+s/2)}{\Gamma(n+1-s/2)}\, r^n \quad \text{for some } C > 0$$

and the right-most series above converges uniformly on compact sets in $[0,1)$ for $n_0 > s/2 > 0$.

Application of the last linear transformations in [1, Eq.s 15.8.1] yields

$$G_n(s;r) = \left(1-r^2\right)^{1-s} \frac{(s/2)_n}{n!} r^n\, {}_2\mathrm{F}_1\left(\begin{matrix} 1-s/2, n+1-s/2\\ n+1\end{matrix}; r^2\right).$$

For s a positive even integer, that is $s = 2m$, we can write (cf. [1, Eq. 15.8.7])

$$G_n(2m;r) = \left(1-r^2\right)^{1-2m} \frac{(m)_n}{n!} \frac{(m)_{m-1}}{(n+1)_{m-1}} r^n\, {}_2\mathrm{F}_1\left(\begin{matrix} 1-m, n+1-m\\ 2-2m\end{matrix}; 1-r^2\right),$$

where the ratios can be simplified further. This shows (9). □

Proof of Proposition 3. The series expansion of the hypergeometric polynomial in (9) yields

$$G_n(2m;r) = \frac{2^{2m-2}\,\Gamma(m-1/2)}{\sqrt{\pi}\,\Gamma(m)} r^n \sum_{k=0}^{m-1} \frac{(1-m)_k (n+1-m)_k}{(2-2m)_k k!} \left(1-r^2\right)^{k+1-2m}.$$

Since

$$\frac{(1-m)_k}{(2-2m)_k} = \binom{m-1}{k} \Big/ \binom{2m-2}{k} = \frac{\Gamma(m)\,\Gamma(2m-1-k)}{\Gamma(m-k)\,\Gamma(2m-1)}$$

and $\Gamma(2m-1) = 2^{2m-2}\,\Gamma(m-1/2)\,\Gamma(m)/\sqrt{\pi}$ (duplication formula for the gamma function) we obtain

$$G_n(2m;r) = \frac{1}{\Gamma(m)} r^n \sum_{k=0}^{m-1} \frac{\Gamma(2m-1-k)}{\Gamma(m-k)k!} \frac{(n+1-m)_k}{k!} \left(1-r^2\right)^{k+1-2m}.$$

By Proposition 2 and the last representation we get

$$\mathcal{M}_{2m}(N;r) = \frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} \left[\sum_{\nu=-\infty}^{\infty} \left(|\nu|N+1-m\right)_k r^{|\nu|N} \right] \left(1-r^2\right)^{k+1-2m} - N\,(1-r)^{-2m}.$$

By Lemma 15 the infinite series reduces to a rational function in r^N. That is

$$\begin{aligned} \mathcal{M}_{2m}(N;r) = \frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} \left[-(1-m)_k + 2\sum_{q=0}^{k} g(k,q;N,1-m)\left(1-r^N\right)^{-q-1} \right] \left(1-r^2\right)^{k+1-2m} \\ - N\,(1-r)^{-2m}, \end{aligned}$$

where $g(k,q;N,b)$ is defined in Lemma 15. $\square$

Proof of Proposition 4. For the Proof of Proposition 4 we need some preparations. We define the following two functions

$$f_q(r) = \left(\frac{1-r^N}{1-r}\right)^{-q}, \qquad h(r) = \frac{1-r^N}{1-r} = \sum_{\ell=0}^{N-1} r^\ell$$

and get a series expansion of $f_q(r)$ at $r=1$. For that we use Faà di Bruno's differentiation formula

$$\{f(g(x))\}^{(n)} = \sum \frac{n!}{k_1!\cdots k_n!} f^{(k)}(g(x)) \prod_{\nu=1}^{n} \left(\frac{g^{(\nu)}(x)}{\nu!}\right)^{k_\nu} = \sum_{k=1}^{n} f^{(k)}(g(x))\,\mathrm{B}_{n,k}(g'(x),g''(x),\dots),$$

where $k = k_1 + \cdots + k_n$ in the first sum and this sum is extended over all partition of n, that is integers $k_1,\dots,k_n \geq 0$ such that $k_1 + 2k_2 + \cdots + nk_n = n$. The polynomials $\mathrm{B}_{n,k}(x_1,x_2,\dots)$ in the second sum are the *(partial) Bell polynomials*, explicitly given by

$$\mathrm{B}_{n,k}(x_1,x_2,\dots) = \sum_{\substack{k_1+k_2+\cdots=k\\ k_1+2k_2+\cdots=n}} \frac{n!}{k_1!k_2!\cdots}\left(\frac{x_1}{1!}\right)^{k_1}\left(\frac{x_2}{2!}\right)^{k_2}\cdots, \tag{15}$$

which satisfy the generating function relation (see, for example, [6])

$$\sum_{n=k}^{\infty} \mathrm{B}_{n,k}(x_1,x_2,x_3,\dots)\frac{t^n}{n!} = \frac{1}{k!}\left(\sum_{m=1}^{\infty} x_m \frac{t^m}{m!}\right)^k. \tag{16}$$

(For the polynomial $\mathrm{B}_{n,k}$ only the variables $x_1, x_2, \dots, x_{n-k+1}$ will be effective.) We record that

$$\mathrm{B}_{n,k}(\alpha x_1, \alpha x_2, \dots) = \alpha^k\, \mathrm{B}_{n,k}(x_1,x_2,\dots), \qquad \alpha \neq 0, \tag{17a}$$

$$\mathrm{B}_{n,k}(\alpha^1 x_1, \alpha^2 x_2, \dots) = \alpha^n\, \mathrm{B}_{n,k}(x_1,x_2,\dots), \qquad \alpha \neq 0. \tag{17b}$$

Lemma 7. *Let n be an integer ≥ 0. Then*

$$h^{(n)}(1) = \frac{N\,(N-1)\cdots(N-n)}{n+1} = n!\binom{N}{n+1}.$$

Proof. One has

$$h^{(n)}(r) = \left\{\frac{1-r^N}{1-r}\right\}^{(n)} = \left\{\sum_{\ell=0}^{N-1} r^\ell\right\}^{(n)} = \sum_{\ell=0}^{N-1}\left\{r^\ell\right\}^{(n)} = \sum_{\ell=n}^{N-1} \ell\,(\ell-1)\cdots(\ell-n)\,r^{\ell-n}$$

with the understanding that a sum with upper index is smaller than the lower index is empty and therefore set to zero. If $r=1$, then

$$h^{(n)}(1) = \sum_{\ell=n}^{N-1} \ell\,(\ell-1)\cdots(\ell-n) = \sum_{\ell=1}^{N-n} (\ell+n-1)\cdots(\ell+1)\,\ell = \sum_{\ell=1}^{N-n} (\ell)_n = \frac{N\,(N-1)\cdots(N-n)}{n+1}.$$

The last step follows by complete induction in N. $\square$

Lemma 8. *Let q be a positive real number and N a positive integer. Then for $m = 1, 2, \dots$*

$$f_q(r) = N^{-q} + \sum_{n=1}^{m} \frac{f_q^{(n)}(1)}{n!}\,(r-1)^n + \mathcal{R}_m(r), \qquad 0 < r < 1,$$

where the coefficients $f_q^{(n)}(1)$ can be written as

$$f_q^{(n)}(1) = \sum_{k=1}^{n} (-1)^k (q)_k N^{-q-k}\; \mathrm{B}_{n,k}(1!\binom{N}{2}, 2!\binom{N}{3}, 3!\binom{N}{4}, \dots)$$

and the remainder can be estimated by

$$|\mathcal{R}_m(r)| \leq \frac{(1-r)^{m+1}}{(m+1)!} \sum_{k=1}^{m+1} (q)_k f_{q+k}(r)\; \mathrm{B}_{m+1,k}(1!\binom{N}{2}, 2!\binom{N}{3}, 3!\binom{N}{4}, \dots).$$

Proof. Let $m \geq 1$. By Taylor's theorem

$$f_q(r) = \left(\frac{1-r^N}{1-r} \right)^{-q} = \sum_{n=0}^{m} \frac{f_q^{(n)}(1)}{n!} \, (r-1)^n + \mathcal{R}_m(r), \qquad \mathcal{R}_m(r) = \frac{f_q^{(m+1)}(\rho)}{(m+1)!} \, (r-1)^{m+1}$$

for some ρ with $r < \rho < 1$. By Faà di Bruno's differentiation formula

$$\begin{aligned} f_q^{(n)}(r) = \left\{ \left(\frac{1-r^N}{1-r} \right)^{-q} \right\}^{(n)} &= \sum_{k=1}^{n} \left\{ (\cdot)^{-q} \right\}^{(k)} (h(r)) \, \mathrm{B}_{n,k}(h'(r), h''(r), \dots) \\ &= \sum_{k=1}^{n} (-1)^k (q)_k \, (h(r))^{-q-k} \, \mathrm{B}_{n,k}(h'(r), h''(r), \dots). \end{aligned}$$

The representation of $f_q^{(n)}(1)$ follows from the fact $h(1) = N$ and Lemma 7.

The remainder $\mathcal{R}_m(r)$ is estimated next. Observe that the function $h^{(\nu)}(r)$ is positive and strictly monotonically increasing on $(0,1)$ for each $\nu \geq 0$. Thus, by the definition of Bell polynomials (15) one gets

$$\begin{aligned} \left| f_q^{(m+1)}(\rho) \right| &\leq \sum_{k=1}^{m+1} (q)_k \, (h(\rho))^{-q-k} \left| \mathrm{B}_{m+1,k}(h'(r), h''(r), \dots) \right| \\ &\leq \sum_{k=1}^{m+1} (q)_k \, (h(r))^{-q-k} \, \mathrm{B}_{m+1,k}(h'(1), h''(1), \dots). \end{aligned}$$

The estimate follows. □

Let $\binom{a_1+\cdots+a_k}{a_1,\dots,a_k}$ denote the multinomial coefficient.

The specific Bell polynomials appearing in Lemma 8 can be represented as follows.

Lemma 9. *Let n, k and N be positive integers. Then*

$$\begin{aligned} \mathrm{B}_{n,k}(1!\binom{N}{2}, 2!\binom{N}{3}, 3!\binom{N}{4}, \dots) &= \frac{n!}{k!} \sum_{\substack{n_1=1 \\ n_1+\cdots+n_k=n}}^{N-1} \cdots \sum_{n_k=1}^{N-1} \binom{N}{n_1+1} \cdots \binom{N}{n_k+1} \\ &= \frac{n!}{k!(n+k)!} \sum_{\ell=k}^{k+n} (-1)^{n-\ell+k} H_\ell(n,k) \, N^\ell, \end{aligned}$$

where the coefficients

$$H_\ell(n,k) = \sum_{\substack{n_1=1 \\ n_1+\cdots+n_k=n}}^{n} \cdots \sum_{n_k=1}^{n} \binom{n+k}{n_1+1, n_2+1, \dots, n_k+1} \sum_{\substack{\ell_1=1 \\ \ell_1+\cdots+\ell_k=\ell}}^{n_1+1} \cdots \sum_{\ell_k=1}^{n_k+1} \left| \mathrm{s}(n_1+1, \ell_1) \right| \cdots \left| \mathrm{s}(n_k+1, \ell_k) \right|.$$

vanish for $\ell = 0, 1, \dots, k-1$ and do not depend on N.

Proof. By the generating function relation for partial Bell polynomials (16)

$$\sum_{n=k}^{\infty} \mathrm{B}_{n,k}(1!\binom{N}{2}, 2!\binom{N}{3}, 3!\binom{N}{4}, \dots) \frac{t^n}{n!} = \frac{1}{k!} \left(\sum_{m=1}^{\infty} \binom{N}{m+1} t^m \right)^k.$$

The right-hand side above evaluates as

$$\begin{aligned}\frac{1}{k!}\left[\frac{(1+t)^N-1-N\,t}{t}\right]^k &= \frac{1}{k!}\left[\sum_{\nu=2}^{N}\binom{N}{\nu}t^{\nu-1}\right]^k = \frac{1}{k!}\left[\sum_{\nu=1}^{N-1}\binom{N}{\nu+1}t^{\nu}\right]^k \\ &= \frac{1}{k!}\sum_{\nu_1=1}^{N-1}\cdots\sum_{\nu_k=1}^{N-1}\binom{N}{\nu_1+1}\cdots\binom{N}{\nu_k+1}t^{\nu_1+\cdots+\nu_k} \\ &= \sum_{\nu=k}^{k(N-1)}\left\{\frac{\nu!}{k!}\sum_{\substack{\nu_1=1 \\ \nu_1+\cdots+\nu_k=\nu}}^{N-1}\cdots\sum_{\nu_k=1}^{N-1}\binom{N}{\nu_1+1}\cdots\binom{N}{\nu_k+1}\right\}\frac{t^{\nu}}{\nu!}.\end{aligned}$$

This shows that the infinite series at the left-hand side above is, in fact, a finite series and comparison of coefficients yields the first identity of our result

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots) = \frac{n!}{k!}\sum_{\substack{n_1=1 \\ n_1+\cdots+n_k=n}}^{N-1}\cdots\sum_{n_k=1}^{N-1}\binom{N}{n_1+1}\cdots\binom{N}{n_k+1}.$$

Let $[x]_n$ denote the *falling factorial* $[x]_n = x(x-1)\cdots(x-n+1)$. Taking into account that $\binom{N}{\nu+1} = [N]_{\nu+1}/(\nu+1)!$ vanishes if $\nu+1>N$ for the positive integer N, we may write

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots) = \frac{n!}{k!}\sum_{\substack{n_1=1 \\ n_1+\cdots+n_k=n}}^{n}\cdots\sum_{n_k=1}^{n}\frac{[N]_{n_1+1}}{(n_1+1)!}\cdots\frac{[N]_{n_k+1}}{(n_k+1)!}$$

By the definition of Stirling numbers of the first kind and using that $n_1+\cdots+n_k=n$ one has

$$\begin{aligned}\frac{[N]_{n_1+1}}{(n_1+1)!}\cdots\frac{[N]_{n_k+1}}{(n_k+1)!} &= \prod_{\nu=1}^{k}\left[\frac{1}{(n_\nu+1)!}\sum_{\ell=0}^{n_\nu+1}s(n_\nu+1,\ell)\,N^{\ell}\right] \\ &= \sum_{\ell_1=0}^{n_1+1}\cdots\sum_{\ell_k=0}^{n_k+1}\frac{s(n_1+1,\ell_1)\cdots s(n_k+1,\ell_k)}{(n_1+1)!\cdots(n_k+1)!}N^{\ell_1+\cdots+\ell_k} \\ &= \sum_{\ell=0}^{k+n}\left\{\sum_{\substack{\ell_1=0 \\ \ell_1+\cdots+\ell_k=\ell}}^{n_1+1}\cdots\sum_{\ell_k=0}^{n_k+1}\frac{s(n_1+1,\ell_1)\cdots s(n_k+1,\ell_k)}{(n_1+1)!\cdots(n_k+1)!}\right\}N^{\ell}.\end{aligned}$$

When expanding the left-hand side above in powers of N we infer that the lowest power appearing at the right-hand side above is N^k. That is

$$\sum_{\substack{\ell_1=0 \\ \ell_1+\cdots+\ell_k=\ell}}^{n_1+1}\cdots\sum_{\ell_k=0}^{n_k+1}\frac{s(n_1+1,\ell_1)\cdots s(n_k+1,\ell_k)}{(n_1+1)!\cdots(n_k+1)!} = 0 \qquad \text{for } \ell=0,1,\dots,k-1.$$

Putting everything together we get the identity

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots) = \frac{n!}{k!}\sum_{\ell=k}^{k+n}\mu_\ell(n,k;N)\,N^{\ell}$$

and the representation

$$\mu_\ell(n,k) = \sum_{\substack{n_1=1 \\ n_1+\cdots+n_k=n}}^{n}\cdots\sum_{n_k=1}^{n}\sum_{\substack{\ell_1=1 \\ \ell_1+\cdots+\ell_k=\ell}}^{n_1+1}\cdots\sum_{\ell_k=1}^{n_k+1}\frac{\mathrm{s}(n_1+1,\ell_1)\cdots\mathrm{s}(n_k+1,\ell_k)}{(n_1+1)!\cdots(n_k+1)!}.$$

It is well-known that $(-1)^{n-k}\,\mathrm{s}(n,k)>0$ for $n\geq k$. Thus

$$\prod_{\nu=1}^{k}\mathrm{s}(n_\nu+1,\ell_\nu)=\prod_{\nu=1}^{k}(-1)^{n_\nu+1-\ell_\nu}\left|\mathrm{s}(n_\nu+1,\ell_\nu)\right|=(-1)^{n_1+\cdots+n_k+k-\ell_1-\cdots-\ell_k}\left|\prod_{\nu=1}^{k}\mathrm{s}(n_\nu+1,\ell_\nu)\right|$$

and from $n_1+\cdots+n_k=n$ and $\ell_1+\cdots+\ell_k=\ell$ it follows that

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots)=\frac{n!}{k!}\sum_{\ell=k}^{k+n}(-1)^{n-\ell+k}\left|\mu_\ell(n,k)\right|N^\ell.$$

By definition $(-1)^{n-k}\,\mathrm{s}(n,k)$ counts the permutations of $1,2,\dots,n$ with exactly k cycles. With this combinatorial interpretation in mind we can write

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots)=\frac{n!}{k!(n+k)!}\sum_{\ell=k}^{k+n}(-1)^{n-\ell+k}H_\ell(n,k)\,N^\ell,$$

where the coefficients $H_\ell(n,k)$ (which do not depend on N) are given in the lemma. □

Special values of the numbers $H_\ell(n,k)$ are as follows.

Lemma 10. *Let n (≥ 1) and k (≥ 1) be integers. Then*

$$H_{n+k}(n,n)=2^{-n}(2n)!\binom{n}{k},$$

$$H_{n+k}(n,k)=\sum_{\substack{n_1=2\\n_1+\cdots+n_k=n+k}}^{n+1}\cdots\sum_{n_k=2}^{n+1}\binom{n+k}{n_1,n_2,\dots,n_k}=\frac{k!(n+k)!}{n!}\,\mathrm{B}_{n,k}(\frac{1}{2},\frac{1}{3},\dots).$$

Proof. By Lemma 9

$$H_{n+k}(n,n)=\sum_{\substack{n_1=1\\n_1+\cdots+n_n=n}}^{n}\cdots\sum_{n_n=1}^{n}\binom{2n}{n_1+1,n_2+1,\dots,n_n+1}\sum_{\substack{\ell_1=1\\\ell_1+\cdots+\ell_n=n+k}}^{n_1+1}\cdots\sum_{\ell_n=1}^{n_n+1}\left|\mathrm{s}(n_1+1,\ell_1)\right|\cdots\left|\mathrm{s}(n_n+1,\ell_n)\right|.$$

It follows that $n_1=\cdots=n_n=1$. Hence, the multinomial coefficient evaluates as $2^{-n}(2n)!$ and

$$H_{n+k}(n,n)=2^{-n}(2n)!\sum_{\substack{\ell_1=0\\\ell_1+\cdots+\ell_n=k}}^{1}\cdots\sum_{\ell_n=0}^{1}\left|\mathrm{s}(2,\ell_1+1)\right|\cdots\left|\mathrm{s}(2,\ell_n+1)\right|.$$

Since $\mathrm{s}(2,1)=-1$ and $\mathrm{s}(2,2)=1$, the sum counts the number of n-tuples $(\ell_1,\dots,\ell_n)$ with precisely k of the entries being 1 which is equal to $\binom{n}{k}$. The result follows.

Similarly

$$H_{n+k}(n,k)=\sum_{\substack{n_1=1\\n_1+\cdots+n_k=n}}^{n}\cdots\sum_{n_k=1}^{n}\binom{n+k}{n_1+1,n_2+1,\dots,n_k+1}\sum_{\substack{\ell_1=1\\\ell_1+\cdots+\ell_k=n+k}}^{n_1+1}\cdots\sum_{\ell_k=1}^{n_k+1}\left|\mathrm{s}(n_1+1,\ell_1)\right|\cdots\left|\mathrm{s}(n_k+1,\ell_k)\right|.$$

Here, the inner sum reduces to a single term with $\ell_1=n_1+1$, …, $\ell_k=n_k+1$ which evaluates to 1. Thus

$$H_{n+k}(n,k)=\sum_{\substack{n_1=1\\n_1+\cdots+n_k=n}}^{n}\cdots\sum_{n_k=1}^{n}\binom{n+k}{n_1+1,n_2+1,\dots,n_k+1}=\sum_{\substack{n_1=2\\n_1+\cdots+n_k=n+k}}^{n+1}\cdots\sum_{n_k=2}^{n+1}\binom{n+k}{n_1,n_2,\dots,n_k}.$$

On the other hand, by Lemma 9

$$\frac{n!}{k!(n+k)!}H_{n+k}(n,k)$$

is the coefficient of the highest power of N in

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots).$$

From

$$\nu!\binom{N}{\nu+1} = \frac{1}{\nu+1}N^{\nu+1}\left(1+\mathcal{O}(N^{-1})\right) \qquad \text{as } N\to\infty$$

and the identities (17) we obtain

$$\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots) = N^{n+k}\,\mathrm{B}_{n,k}(\frac{1+\mathcal{O}(N^{-1})}{2},\frac{1+\mathcal{O}(N^{-1})}{3},\dots) \qquad \text{as } N\to\infty.$$

It follows that

$$\frac{n!}{k!(n+k)!}H_{n+k}(n,k) = \mathrm{B}_{n,k}(\frac{1}{2},\frac{1}{3},\dots), \qquad n\geq k\geq 1.$$

□

Lemma 9 allows us to recast the representation of $f_q^{(n)}(1)$ given in Lemma 8.

Corollary 11. *Let q, n and N be positive integer. Then*

$$\frac{f_q^{(n)}(1)}{n!} = \sum_{\ell=0}^{n}(-1)^\ell G(n,\ell,q)\,N^{\ell-q}, \qquad G(n,\ell,q) = \sum_{k=1}^{n}\frac{(-1)^{n-k}(q)_k}{k!(n+k)!}H_{\ell+k}(n,k),$$

where the numbers $H_{\ell+k}(n,k)$ are given in Lemma 9.

Proof. From Lemmas 8 and 9

$$\begin{aligned}\frac{f_q^{(n)}(1)}{n!} &= \frac{1}{n!}\sum_{k=1}^{n}(-1)^k(q)_kN^{-q-k}\,\mathrm{B}_{n,k}(1!\binom{N}{2},2!\binom{N}{3},3!\binom{N}{4},\dots)\\ &= \frac{1}{n!}\sum_{k=1}^{n}(-1)^k(q)_kN^{-q-k}\,\frac{n!}{k!(n+k)!}\sum_{\ell=k}^{k+n}(-1)^{n-\ell+k}H_\ell(n,k)\,N^\ell\\ &= \sum_{k=1}^{n}(-1)^k(q)_kN^{-q-k}\,\frac{1}{k!(n+k)!}\sum_{\ell=0}^{n}(-1)^{n-\ell}H_{\ell+k}(n,k)\,N^{\ell+k}\\ &= \sum_{\ell=0}^{n}(-1)^\ell\sum_{k=1}^{n}\frac{(-1)^{n-k}(q)_k}{k!(n+k)!}H_{\ell+k}(n,k)\,N^{\ell-q}.\end{aligned}$$

The result follows. □

Lemma 12. *Let q be a positive integer. Then one has the generating function relation*

$$\sum_{n=1}^{\infty}(-1)^nG(n,n,q)\,x^n = \left(\frac{e^x-1}{x}\right)^{-q} - 1.$$

Proof. By Corollary 11 and Lemma 10

$$(-1)^nG(n,n,q) = \sum_{k=1}^{n}\frac{(-1)^k(q)_k}{k!(n+k)!}H_{n+k}(n,k) = \frac{1}{n!}\sum_{k=1}^{n}(-1)^k(q)_k\,\mathrm{B}_{n,k}(1/2,1/3,\dots).$$

Application of the identities (cf. Faà di Bruno's differentiation formula)

$$g(f(x)) = \sum_{n=1}^{\infty}\frac{\sum_{k=1}^{n}b_k\,\mathrm{B}_{n,k}(a_1,a_2,\dots)}{n!}x^n, \qquad f(x) = \sum_{n=1}^{\infty}\frac{a_n}{n!}x^n, \quad g(y) = \sum_{n=1}^{\infty}\frac{b_n}{n!}y^n,$$

to the functions

$$f(x) = \sum_{n=1}^{\infty}\frac{x^n}{(n+1)!} = \frac{e^x-1-x}{x}, \qquad g(y) = \sum_{n=1}^{\infty}\frac{(-1)^n(q)_n}{n!}y^n = (1+y)^{-q}-1$$

yields

$$\sum_{n=1}^{\infty}(-1)^nG(n,n,q)\,x^n = g(f(x)) = \left(\frac{e^x-1}{x}\right)^{-q} - 1.$$

□

With these preparations we are ready to give the following proof.

Proof of Proposition 4. The formula of $\mathcal{M}_{2m}(N;r)$ in Prop. 3 is rearranged w.r.t. powers of $(1-r)$. That is

$$\begin{aligned}\mathcal{M}_{2m}(N;r) + N\left(1-r\right)^{-2m} &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} (1-m)_k \left(1+r\right)^{k+1-2m} \left(1-r\right)^{k+1-2m} \\ &+ 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} g(k,q;N,1-m) \left(\frac{1-r^N}{1-r}\right)^{-q-1} \left(1+r\right)^{k+1-2m} \left(1-r\right)^{k-q-2m}.\end{aligned}$$

Note that $k+1-2m<0$. Using Taylor expansion (cf. Lemma 8, since $f_q(r) = (1+r)^{-q}$ if $N=2$)

$$\left(1+r\right)^{-q} = \sum_{n=0}^{m} \frac{(q)_n}{n!} 2^{-q-n} \left(1-r\right)^n + \tilde{\mathcal{R}}_m(r), \qquad \left|\tilde{\mathcal{R}}_m(r)\right| \leq \frac{(q)_{m+1}}{(m+1)!} \left(1-r\right)^{m+1},$$

one gets (N is assumed to be fixed)

$$\begin{aligned}\mathcal{M}_{2m}(N;r) + N\left(1-r\right)^{-2m} &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} (1-m)_k \\ &\quad\times \sum_{n=0}^{2m-1-k} \frac{(2m-1-k)_n}{n!} 2^{k-n+1-2m} \left(1-r\right)^{k+n+1-2m} + \mathcal{O}((1-r)) \\ &+ 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} g(k,q;N,1-m) \left(\frac{1-r^N}{1-r}\right)^{-q-1} \\ &\quad\times \sum_{n=0}^{2m+q-k} \frac{(2m-1-k)_n}{n!} 2^{k-n+1-2m} \left(1-r\right)^{n+k-q-2m} + \mathcal{O}((1-r)) \\ &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{n=0}^{2m-1-k} \frac{\Gamma(n+2m-k-1)}{\Gamma(m-k)k!n!} (1-m)_k 2^{k+1-2m-n} \left(1-r\right)^{n+k+1-2m} \\ &+ 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \frac{\Gamma(n+2m-k-1)}{\Gamma(m-k)k!n!} g(k,q;N,1-m) \left(\frac{1-r^N}{1-r}\right)^{-q-1} 2^{k+1-2m-n} \left(1-r\right)^{n+k-q-2m} \\ &+ \mathcal{O}((1-r)), \qquad \text{as } r \to 1^-.\end{aligned}$$

By Lemma 8

$$\begin{aligned}\mathcal{M}_{2m}(N;r) + N\left(1-r\right)^{-2m} &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{n=0}^{2m-1-k} \frac{\Gamma(n+2m-k-1)}{\Gamma(m-k)k!n!} (1-m)_k 2^{k+1-2m-n} \left(1-r\right)^{n+k+1-2m} \\ &+ 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \frac{\Gamma(n+2m-k-1)}{\Gamma(m-k)k!n!} g(k,q;N,1-m) N^{-1-q} 2^{k+1-2m-n} \left(1-r\right)^{n+k-q-2m} \\ &+ 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \sum_{\nu=1}^{2m+q-k-n} \frac{\Gamma(n+2m-k-1)}{\Gamma(m-k)k!n!} g(k,q;N,1-m) \frac{(-1)^\nu f_{q+1}^{(\nu)}(1)}{\nu!} \\ &\quad\times 2^{k+1-2m-n} \left(1-r\right)^{\nu+n+k-q-2m} + \mathcal{O}((1-r)), \qquad \text{as } r \to 1^-.\end{aligned}$$

The power $(1-r)^{-2m}$ appears only in the triple sum when $q=k$ and $n=0$. Its coefficient is

$$c_{-2m} {:=} 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m-k)k!} g(k,k;N,1-m) N^{-1-k} 2^{k+1-2m}.$$

By the formulas in Prop. 3 and the properties of the Stirling and Eulerian numbers (cf. (23))

$$g(k,k;N,1-m) = \mathrm{s}(k,k;1-m)b(k,0)\,N^k = s(k,k)\,N^k \sum_{j=0}^{k} \left\langle {k \atop j} \right\rangle = s(k,k)k!N^k = k!N^k.$$

Thus, by [8, Eq. 4.2.3.10],

$$c_{-2m} = N2^{2-2m} \sum_{k=0}^{m-1} \frac{\Gamma(2m-k-1)}{\Gamma(m)\,\Gamma(m-k)} 2^k = N2^{2-2m} \sum_{k=0}^{m-1} \binom{2m-2-k}{m-1} 2^k = N$$

and, therefore, the term $N(1-r)^{-2m}$ could be removed from both sides of the identity above. First, we reorder the terms

$$\begin{aligned}
&\mathcal{M}_{2m}(N;r) + N\,(1-r)^{-2m} = -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{n=0}^{2m-1-k} \frac{\Gamma(2(2m-1-k)-n)}{\Gamma(m-k)k!(2m-1-k-n)!} (1-m)_k 2^{n-2(2m-1-k)}\,(1-r)^{-n} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} g(k,q;N,1-m) N^{-1-q} 2^{n-q-1-2(2m-1-k)}\,(1-r)^{-n} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \sum_{\nu=1}^{n} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} g(k,q;N,1-m) \frac{(-1)^\nu f_{q+1}^{(\nu)}(1)}{\nu!} \\
&\qquad \times 2^{n-q-1-2(2m-1-k)}\,(1-r)^{\nu-n} + \mathcal{O}((1-r)) \\
&= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{n=0}^{m+k} \frac{\Gamma(2(m+k)-n)}{\Gamma(m-k)k!(m+k-n)!} (1-m)_{m-1-k} 2^{n-2(m+k)}\,(1-r)^{-n} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} g(k,q;N,1-m) N^{-1-q} 2^{n-q-1-2(2m-1-k)}\,(1-r)^{-n} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \sum_{\nu=0}^{n-1} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} g(k,q;N,1-m) \frac{(-1)^{n-\nu} f_{q+1}^{(n-\nu)}(1)}{(n-\nu)!} \\
&\qquad \times 2^{n-q-1-2(2m-1-k)}\,(1-r)^{-\nu} + \mathcal{O}((1-r)), \qquad \text{as } r \to 1^-.
\end{aligned}$$

Since $\lim_{r\to 1^-} \mathcal{M}_{2m}(N;r) = \mathcal{M}_{2m}(N)$ and $\mathcal{M}_{2m}(N)$ exists and is finite for all positive integers m and N with $N \geq 2$ (and $N(1-r)^{-2m}$ can be removed from both sides of the identities above) it follows that the cumulative coefficient of a negative power of $(1-r)$ is zero and we are left with

$$\begin{aligned}
\mathcal{M}_{2m}(N) &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2(m+k))}{\Gamma(m-k)k!(m+k)!} (1-m)_{m-1-k} 2^{-2(m+k)} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \frac{\Gamma(2(2m-1-k)+q+1)}{\Gamma(m-k)k!(2m+q-k)!} g(k,q;N,1-m) N^{-1-q} 2^{-q-1-2(2m-1-k)} \\
&\quad + 2\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} g(k,q;N,1-m) \frac{(-1)^n f_{q+1}^{(n)}(1)}{n!} 2^{n-q-1-2(2m-1-k)}.
\end{aligned}$$

Substituting the expressions for $g(k,q;N,1-m)$ (cf. Prop. 3) and $f_{q+1}^{(n)}(1)$ (cf. Cor. 11) we get

$$\begin{aligned}
\mathcal{M}_{2m}(N) &= -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2(m+k))}{\Gamma(m-k)k!(m+k)!} (1-m)_{m-1-k} 2^{-2(m+k)} \\
&\quad + N \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{p=0}^{k-q} X(k,q,p;m) N^p + N \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \sum_{p=0}^{k-q} \sum_{\ell=0}^{n} X(k,q,p,n,\ell;m)\,N^{\ell+p},
\end{aligned} \tag{18}$$

where the primary coefficients are given by

$$\begin{aligned}
X(k,q,p;m) &:= \frac{2}{\Gamma(m)} \frac{\Gamma(2(2m-1-k)+q+1)}{\Gamma(m-k)k!(2m+q-k)!} \frac{(-1)^p s(k,p+q;1-m) b(p+q,p)}{2^{q+1+2(2m-1-k)}} \\
X(k,q,p,n,\ell;m) &:= \frac{2}{\Gamma(m)} \frac{\Gamma(2(2m-1-k)+q-n+1)}{\Gamma(m-k)k!(2m+q-k-n)!} \frac{(-1)^{p+n+\ell} s(k,p+q;1-m) b(p+q,p)}{2^{q+1+2(2m-1-k)-n}} G(n,\ell,q+1)
\end{aligned}$$

which depend on the secondary coefficients

$$\mathrm{s}(n,\ell;y) = \sum_{k=\ell}^{n} \binom{k}{\ell} \mathrm{s}(n,k) \left(y+n-1\right)^{k-\ell}, \qquad b(p+q,p) = \sum_{j=0}^{q} \left\langle {p+q \atop j} \right\rangle \binom{p+q-j}{p}$$

and $G(n,\ell,q)$ is given in Cor. 11 (cf. Lemma 9).

We reorder the terms in (18) w.r.t. powers of N. The first part is

$$\sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{p=0}^{k-q} X(k,q,p;m) N^p = \sum_{k=0}^{m-1} \sum_{p=0}^{k} \sum_{q=0}^{k-p} X(k,q,p;m) N^p = \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} \sum_{q=0}^{k-p} X(k,q,p;m) N^p.$$

The second part is

$$\begin{aligned}
&\sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{n=0}^{2m+q-k} \sum_{p=0}^{k-q} \sum_{\ell=0}^{n} X(k,q,p,n,\ell;m)\, N^{\ell+p} = \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{p=0}^{q} \sum_{n=0}^{2m-q} \sum_{\ell=0}^{n} X(k,k-q,p,n,\ell;m)\, N^{\ell+p} \\
&\quad = \sum_{k=0}^{m-1} \sum_{q=0}^{k} \sum_{p=0}^{q} \sum_{\ell=0}^{2m-q} \sum_{n=\ell}^{2m-q} X(k,k-q,p,n,\ell;m)\, N^{\ell+p} = \sum_{k=0}^{m-1} \sum_{p=0}^{k} \sum_{q=p}^{k} \sum_{\ell=0}^{2m-q} \sum_{n=\ell}^{2m-q} X(k,k-q,p,n,\ell;m)\, N^{\ell+p} \\
&\quad = \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} \sum_{q=p}^{k} \sum_{\ell=0}^{2m-q} \sum_{n=\ell}^{2m-q} X(k,k-q,p,n,\ell;m)\, N^{\ell+p} = \sum_{p=0}^{m-1} \sum_{q=p}^{m-1} \sum_{\ell=0}^{2m-q} \sum_{k=q}^{m-1} \sum_{n=\ell}^{2m-q} X(k,k-q,p,n,\ell;m)\, N^{\ell+p} \\
&\quad = \left[\sum_{p=0}^{m-1} \sum_{\ell=0}^{m} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\ell}^{2m-q} + \sum_{p=0}^{m-1} \sum_{\ell=m+1}^{2m-p} \sum_{q=p}^{2m-\ell} \sum_{k=q}^{m-1} \sum_{n=\ell}^{2m-q} \right] X(k,k-q,p,n,\ell;m)\, N^{\ell+p} \\
&\quad = \left[\sum_{\nu=0}^{m-1} \sum_{p=0}^{\nu} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} + \sum_{\nu=m}^{2m-1} \sum_{p=\nu-m}^{m-1} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} \right] X(k,k-q,p,n,\nu-p;m)\, N^{\nu} \\
&\qquad + \sum_{\nu=m+1}^{2m} \sum_{p=0}^{\nu-m-1} \sum_{q=p}^{2m-\nu+p} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} X(k,k-q,p,n,\nu-p;m)\, N^{\nu}.
\end{aligned}$$

Putting everything together we arrive at

$$\begin{aligned}
\mathcal{M}_{2m}(N) = -\frac{N^2}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2(m+k))}{\Gamma(m-k)k!(m+k)!} (1-m)_{m-1-k} 2^{-2(m+k)} + N \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} \sum_{q=0}^{k-p} X(k,q,p;m) N^p \\
+ N \left[\sum_{\nu=0}^{m-1} \sum_{p=0}^{\nu} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} + \sum_{\nu=m}^{2m-1} \sum_{p=\nu-m}^{m-1} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} \right] X(k,k-q,p,n,\nu-p;m)\, N^{\nu} \\
+ N \sum_{\nu=m+1}^{2m} \sum_{p=0}^{\nu-m-1} \sum_{q=p}^{2m-\nu+p} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} X(k,k-q,p,n,\nu-p;m)\, N^{\nu}.
\end{aligned}$$

Direct computations (with the help of Mathematica) give the expected results (cf. (5))

$$\begin{aligned}
\mathcal{M}_2(N) &= \frac{N^3}{12} - \frac{N}{12}, \\
\mathcal{M}_4(N) &= -\frac{N^2}{8} - N\left(\frac{N}{4} - \frac{3}{8}\right) + N\left(\frac{3}{8}N - \frac{281}{720}\right) - N\left(\frac{N^3}{16} - \frac{N^2}{72}\right) + N\left(\frac{N^4}{720} + \frac{N^3}{16}\right) \\
&= \frac{N^5}{720} + \frac{10}{720}N^3 - \frac{11}{720}N.
\end{aligned}$$

In general, one has for $m \geq 2$

$$\mathcal{M}_{2m}(N) = \sum_{\nu=0}^{2m} \beta_\nu(m)\, N^{1+\nu},$$

where for $\nu = 0, 1, \dots, m-1$ and $\nu \neq 1$

$$\beta_\nu(m) := \sum_{k=\nu}^{m-1} \sum_{q=0}^{k-\nu} X(k,q,\nu;m) + \sum_{p=0}^{\nu} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} X(k,k-q,p,n,\nu-p;m), \tag{19a}$$

for $\nu = m$ and $m \neq 1$

$$\beta_m(m) := \sum_{p=0}^{m-1} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=m-p}^{2m-q} X(k,k-q,p,n,m-p;m), \tag{19b}$$

for $\nu = m+1, \dots, 2m-1$ (and $m \geq 2$)

$$\begin{aligned}
\beta_\nu(m) :=& \sum_{p=\nu-m}^{m-1} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} X(k,k-q,p,n,\nu-p;m) \\
&+ \sum_{p=0}^{\nu-m-1} \sum_{q=p}^{2m-\nu+p} \sum_{k=q}^{m-1} \sum_{n=\nu-p}^{2m-q} X(k,k-q,p,n,\nu-p;m),
\end{aligned} \tag{19c}$$

for $\nu = 2m$ (and $m \geq 2$)

$$\beta_{2m}(m) := \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} X(k,k-p,p,2m-p,2m-p;m) \tag{19d}$$

and for $\nu = 1$ (and $m \geq 2$)

$$\begin{aligned}
\beta_1(m) =& -\frac{1}{\Gamma(m)} \sum_{k=0}^{m-1} \frac{\Gamma(2(m+k))}{\Gamma(m-k)k!(m+k)!} (1-m)_{m-1-k} 2^{-2(m+k)} \\
&+ \sum_{k=1}^{m-1} \sum_{q=0}^{k-1} X(k,q,1;m) + \sum_{p=0}^{1} \sum_{q=p}^{m-1} \sum_{k=q}^{m-1} \sum_{n=1-p}^{2m-q} X(k,k-q,p,n,1-p;m).
\end{aligned} \tag{19e}$$

The result follows. □

Next the coefficients $\beta_\nu(m)$ are studied in more detail.

Proof of Proposition 5. By the definition of the numbers $X(k,q,p,n,\ell;m)$ given in the Proof of Prop. 4 we get

$$\begin{aligned}
\beta_{2m}(m) &= \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} X(k,k-p,p,2m-p,2m-p;m) \\
&= \sum_{p=0}^{m-1} \sum_{k=p}^{m-1} (-1)^p \frac{2^{2+k-2m}\,\Gamma(2m-1-k)}{\Gamma(m)k!\,\Gamma(m-k)} \,\mathrm{s}(k,k;1-m) b(k,p) G(2m-p,2m-p,k-p+1).
\end{aligned}$$

By the definitions of $\mathrm{s}(n,k;x)$ (Lemma 13) and $b(n,\ell)$ (Lemma 14) we have

$$\mathrm{s}(k,k;1-m) = \mathrm{s}(k,k) = 1, \qquad b(k,p) = \sum_{j=0}^{k-p} \left\langle {k \atop j} \right\rangle \binom{k-j}{p}$$

and the numbers $G(n,n,q)$ are defined by the generating function relation (Lemma 12)

$$\sum_{n=1}^{\infty} (-1)^n G(n,n,q)\, x^n = \left(\frac{e^x - 1}{x} \right)^{-q} - 1.$$

The result follows after interchanging the two summation signs. □

Acknowledgement: The research was supported, in part, by an APART-Fellowship of the Austrian Academy of Sciences. The author is grateful for the hospitality of School of Mathematics and Statistics at University of New South Wales where this research was conducted.

Appendix A. Auxiliary results

Lemma 13. *Let n be a nonnegative integer. Then*

$$(x+y)_n = \sum_{\ell=0}^{n} \mathrm{s}(n,\ell;y)\, x^\ell, \qquad \mathrm{s}(n,\ell;y) := \sum_{k=\ell}^{n} \binom{k}{\ell} \mathrm{s}(n,k)\, (y+n-1)^{k-\ell}.$$

Proof. Using (10) and the Binomial theorem, one gets

$$\begin{aligned}(x+y)_n &= ((x+y+n-1)+(1-n))_n = \sum_{k=0}^{n} \mathrm{s}(n,k)\, (x+y+n-1)^k \\ &= \sum_{k=0}^{n} \sum_{\ell=0}^{k} \mathrm{s}(n,k) \binom{k}{\ell} x^\ell\, (y+n-1)^{k-\ell}.\end{aligned}$$

The result follows after reordering the sum. □

Note that

$$\mathrm{s}(n,0;y) = \sum_{k=0}^{n} \mathrm{s}(n,k)\, (y+n-1)^k = (y+n-1+1-n)_n = (y)_n. \tag{20}$$

Alternatively, one can use the identities ($\left[{n \atop k} \right]$ is the unsigned Stirling number of the first kind)

$$(a+b)_n = \sum_{j=0}^{\infty} \binom{n}{j} (a)_{n-j} (b)_j, \qquad (x)_n = \sum_{k=0}^{n} \left[{n \atop k} \right] x^k$$

to obtain the representation

$$\mathrm{s}(n,\ell;y) = \sum_{k=\ell}^{n} \binom{n}{k} \left[{k \atop \ell} \right] (y)_{n-k}.$$

We need the *polylogarithm function* defined by

$$\mathrm{L}_\nu(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^\nu}, \qquad |z| < 1. \tag{21}$$

In particular, one has for $\nu = -n$ and n a positive integer the relations (cf. [7], [9])

$$\mathrm{L}_{-n}(z) = \sum_{k=1}^{\infty} k^n z^k = \frac{1}{(1-z)^{n+1}} \sum_{j=0}^{n-1} \left\langle {n \atop j} \right\rangle z^{n-j}, \qquad |z| < 1. \tag{22}$$

Lemma 14. *Let n be a positive integer. Then*

$$\mathrm{L}_{-n}(z) = \frac{n! + \sum_{\ell=1}^{n} (-1)^\ell b(n,\ell) \, (1-z)^\ell}{(1-z)^{n+1}}, \qquad b(n,\ell) {:=} \sum_{j=0}^{n-\ell} \left\langle {n \atop j} \right\rangle \binom{n-j}{\ell}.$$

Proof. By (22) and the Binomial theorem

$$\mathrm{L}_{-n}(z) = \frac{1}{(1-z)^{n+1}} \sum_{j=0}^{n-1} \left\langle {n \atop j} \right\rangle [1-(1-z)]^{n-j} = \frac{1}{(1-z)^{n+1}} \sum_{j=0}^{n-1} \sum_{\ell=0}^{n-j} \left\langle {n \atop j} \right\rangle \binom{n-j}{\ell} (-1)^\ell \, (1-z)^\ell .$$

Reordering of the sum yields

$$\mathrm{L}_{-n}(z) = \frac{1}{(1-z)^{n+1}} \left\{ \sum_{j=0}^{n-1} \left\langle {n \atop j} \right\rangle + \sum_{\ell=1}^{n} \left[\sum_{j=0}^{n-\ell} \left\langle {n \atop j} \right\rangle \binom{n-j}{\ell} \right] (-1)^\ell \, (1-z)^\ell \right\}$$

The sum over the Eulerian numbers gives $n!$ (cf. [1, Eq. 26.14.10]). The result follows. □

Note that (using $\left\langle {q \atop q} \right\rangle = 0$ for $q \geq 1$)

$$b(n,0) = \sum_{j=0}^{n} \left\langle {n \atop j} \right\rangle = \sum_{j=0}^{n-1} \left\langle {n \atop j} \right\rangle = n!. \tag{23}$$

Lemma 15. *Let k be a non-negative integer and a, b complex numbers. Then*

$$\sum_{\nu=-\infty}^{\infty} (|\nu| a + b)_k z^{|\nu|} = -(b)_k + 2 \sum_{q=0}^{k} g(k,q;a,b) \, (1-z)^{-q-1}, \qquad |z| < 1,$$

where ($s(n,\ell;y)$ is defined in Lemma 13 and $b(n,\ell)$ is defined in Lemma 14)

$$g(k,q;a,b) {:=} \sum_{p=0}^{k-q} (-1)^p s(k,p+q;b) b(p+q,p) a^{p+q}.$$

Proof. Let $f(z)$ denote the inifinte series above. For $k=0$ one gets

$$f(z) = \sum_{\nu=-\infty}^{\infty} z^{|\nu|} = 1 + 2 \sum_{\nu=1}^{\infty} z^\nu = 1 + \frac{2z}{1-z} = \frac{1+z}{1-z}.$$

Let $k \geq 1$. By Lemma 13 one has

$$f(z) = (b)_k + 2 \sum_{\nu=1}^{\infty} (\nu a + b)_k z^\nu = (b)_k + 2 \sum_{p=0}^{k} \mathrm{s}(k,p;b) a^p \sum_{\nu=1}^{\infty} \nu^p z^\nu = (b)_k + 2 \sum_{p=0}^{k} \mathrm{s}(k,p;b) a^p \, \mathrm{L}_{-p}(z).$$

By Lemma 14 and (10) (and using Eq.s (20) and (23)) one gets

$$\begin{aligned} f(z) &= (b)_k + 2 \, \mathrm{s}(k,0;b) \, \mathrm{L}_0(z) + 2 \sum_{p=1}^{k} \mathrm{s}(k,p;b) a^p \frac{p! + \sum_{\ell=1}^{p} (-1)^\ell b(p,\ell) \, (1-z)^\ell}{(1-z)^{p+1}} = (b)_k + 2 (b)_k \frac{z}{1-z} \\ &\quad + 2 \sum_{p=1}^{k} \mathrm{s}(k,p;b) p! a^p \, (1-z)^{-p-1} + 2 \sum_{p=1}^{k} \sum_{\ell=1}^{p} \mathrm{s}(k,p;b) a^p (-1)^\ell b(p,\ell) \, (1-z)^{-(p+1-\ell)} . \end{aligned}$$

Simplification of the first three terms and summing up over equal powers of $(1-z)$ in the double sum yields

$$\begin{aligned}
f(z) &= -(b)_k + 2\sum_{p=0}^{k} \mathrm{s}(k,p;b)p!a^p\,(1-z)^{-p-1} + 2\sum_{q=0}^{k-1}\left[\sum_{p=1+q}^{k} s(k,p;b)a^p(-1)^{p-q}b(p,p-q)\right](1-z)^{-q-1} \\
&= -(b)_k + 2\,\mathrm{s}(k,k;b)k!a^k\,(1-z)^{-k-1} + 2\sum_{q=0}^{k-1}\left[\mathrm{s}(k,q;b)q!a^q + \sum_{p=1+q}^{k} s(k,p;b)a^p(-1)^{p-q}b(p,p-q)\right](1-z)^{-q-1} \\
&= -(b)_k + 2\,\mathrm{s}(k,k;b)k!a^k\,(1-z)^{-k-1} + 2\sum_{q=0}^{k-1}\left[\sum_{p=q}^{k} s(k,p;b)a^p(-1)^{p-q}b(p,p-q)\right](1-z)^{-q-1} \\
&= -(b)_k + 2\sum_{q=0}^{k}\left[\sum_{p=q}^{k} s(k,p;b)a^p(-1)^{p-q}b(p,p-q)\right](1-z)^{-q-1} \\
&= -(b)_k + 2\sum_{q=0}^{k}\left[\sum_{p=0}^{k-q}(-1)^p s(k,p+q;b)b(p+q,p)a^{p+q}\right](1-z)^{-q-1},
\end{aligned}$$

where we used that $\left\langle {n \atop n} \right\rangle = 0$ for $n \geq 1$ and (23). □

REFERENCES

[1] Digital Library of Mathematical Functions. 2010-05-07. National Institute of Standards and Technology from http://dlmf.nist.gov/.

[2] M. Abramowitz and I. A. Stegun, editors. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. Dover Publications Inc., New York, 1992. Reprint of the 1972 edition.

[3] G. Björck. Distributions of positive mass, which maximize a certain generalized energy integral. *Ark. Mat.*, 3:255–269, 1956.

[4] J. S. Brauchart, D. P. Hardin, and E. B. Saff. The Riesz energy of the Nth roots of unity: an asymptotic expansion for large N. *Bull. Lond. Math. Soc.*, 41(4):621–633, 2009.

[5] W.-C. Chu. Summations on trigonometric functions. *Appl. Math. Comput.*, 141(1):161–176, 2003. Advanced special functions and related topics in differential equations (Melfi, 2001).

[6] M. Mihoubi. Partial bell polynomials and inverse relations. *J. Integer Seq.*, 13(4):Article 10.4.5, 8, 2010.

[7] J. Milnor. On polylogarithms, Hurwitz zeta functions, and the Kubert identities. *Enseign. Math. (2)*, 29(3-4):281–322, 1983.

[8] A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev. *Integrals and series. Vol. 1*. Gordon & Breach Science Publishers, New York, 1986. Elementary functions, Translated from the Russian and with a preface by N. M. Queen.

[9] E. W. Weisstein. Dirichlet L-Series. From MathWorld–A Wolfram Web Resource. http://mathworld.wolfram.com/Polylogarithm.html, Nov. 29 2010.

J. S. BRAUCHART: SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY, NSW, 2052, AUSTRALIA

E-mail address: `j.brauchart@unsw.edu.au`